

## Literatuur artikel bioturbatie

- Breukelaar, A.W., Lammens, E.H.R.R., Klein Breteler, J.G.P., Tátrai, I. (1994). Effects of benthivorous bream (*Abramis brama*) and carp (*Cyprinus carpio*) on sediment resuspension and concentrations of nutrients and chlorophyll a. Freshwater Biology vol. 32, p. 113-121.
- De Lucas Pardo, M.A., Bakker, M., van Kessel ,T., Cozzoli, F., Winterwerp, J.C. (2013.) Erodibility of soft freshwater sediments in Markermeer: the role of bioturbation by meiobenthic fauna. Ocean Dynamics vol. 63, p 1137-1150.
- Gulatie, R.D. & van Donk, E. (2002). Lakes in the Netherlands: their origin, eutrophication and restoration: state of the art review. Hydrobiologia no 478, p. 73–106.
- Hofmann, H., Lorke, A., Peeters , F. (2008). The relative importance of wind and ship waves in the littoral zone of a large lake Limnol. Oceanogr.vol. 53, no. 1, p. 368–380.
- Hurdle, D.P., Stive, R.J.H. (1989). Revision of SPM 1984 Wave Hindcast Model to Avoid Inconsistencies in Engineering Applications. Coastal Engineering vol. 12, p. 339-351.
- Soulsby, R.L., Hamm, L., Klopman, G., Myrhaug, D., Simons, R.R. and Thomas, G.P. (1993). Wave-current interaction within and outside the bottom boundary layer, Coastal Engineering, vol. 21, p. 41-69.
- Tarvainen, M., Ventelä, A.M., Helminen, H., Sarvala, J. (2005) Nutrient release and resuspension generated by ruffe (*Gymnocephalus cernuus*) and chironomids. Freshwater Biology vol. 50, p. 447-458.

